

REMARK 1. The above approach of this paper can be combined with the acceleration of the bisection eigenvalue algorithm proposed in [4] and based on the techniques that ensure superlinear (quadratic to cubic) convergence (right from the start) to the isolated eigenvalues or isolated clusters of k eigenvalues based on the Newton-like formula $x_{i+1} = x_i - k p(x_i)/p'(x_i)$, $i = 0, 1, \dots$, and on its refinements.

REMARK 2. Bini and Pan [2] use a fast (but generally numerically unstable) algorithm for simultaneous multipoint polynomial evaluation, in order to accelerate the dc algorithm; Gu and Eisenstat [5] propose to accelerate the dc algorithm by using the multipole algorithm for approximating the secular (rational) function involved.

REMARK 3. A unitary matrix U and the Hermitian matrix $U + U^H$ have the same eigenvectors; moreover, the sets of the eigenvalues of U and of $U + U^H$ are easily expressed through each other. Thus, the above approach can be extended to the eigenproblem for unitary matrices.

For the unsymmetric eigenvalue problem, the bisection method can be extended in the form of Weyl's construction, known for approximating polynomial zeros [6,7] and consisting of recursive proximity tests. Such a test computes, within the relative error $\sqrt{2} - 1$, the distance from the center of a square on the complex plane to the nearest zero of a given polynomial (in our case, to the nearest eigenvalue of a matrix A). If the test proves that a square is eigenvalue-free, the square is discarded. The test (for the eigenvalues) can be performed probabilistically, at the cost of $O(n^2 \log n)$ flops, for a Hessenberg matrix A [8]. The resulting algorithm is as robust as the bisection method and uses $O(bn^3 \log n)$ flops to approximate all (the clusters of) the eigenvalues of A within the error bound $2^{-b}\|A\|_1$. Numerical contour integration can then be applied to count the multiplicity of the eigenvalues or their numbers in each of their clusters.

REFERENCES

1. V. Pan, Linear time approximate evaluation of a polynomial at real points, Tech. Report TR 92-4, Comp. Sci. Dept. SUNYA, Albany, NY, (1992).
2. D. Bini and V. Pan, Parallel complexity of tridiagonal symmetric eigenvalue problem, In *Proc. 2nd Ann. ACM-SIAM Symp. on Discrete Algorithms*, pp. 384-393, (1991).
3. D. Bini and V. Pan, Practical improvement of the divide-and-conquer eigenvalue algorithms, *Computing* **48**, 109-123 (1992).
4. V. Pan, Accelerated solution of the symmetric eigenvalue problem, Tech. Report TR 92-6, Computer Science Dept., SUNYA, Albany, NY, (1992).
5. M. Gu and S.C. Eisenstat, private communication.
6. P. Henrici, *Applied and Computational Complex Analysis*, John Wiley, New York, (1974).
7. V. Pan, Sequential and parallel complexity of approximate evaluation of polynomial zeros, *Computers & Math. with Applications* **14** (8), 591-622 (1987).
8. J.D. Dixon, Estimating extremal eigenvalues and condition numbers of matrices, *SIAM J. of Numer. Analysis* **20** (4), 812-814 (1983).